

# Soil CO<sub>2</sub> evolution from Korean pine virgin forest at Changbai Mountain<sup>1</sup>

Ma Yueqiang (马越强) Yan Xiaodong (延晓冬) Yang Sihe (杨思河)

Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110015, P. R. China

**Abstract** The soil CO<sub>2</sub> evolution rate was measured in a virgin Korean pine forest. The results in June showed that the lowest value of evolution rate was 220 mg/(m<sup>2</sup>·h) and appeared at 6:00 a.m. The highest value was 460 mg/(m<sup>2</sup>·h) at 18:00. The rates of CO<sub>2</sub> evolution were related with soil temperature. On the basis of the constructed regression equation and the monthly average values of temperature, the magnitude of CO<sub>2</sub> evolution from Korean pine forest soil was 10.4 t/hm<sup>2</sup> during a growing season.

**Key words:** Soil CO<sub>2</sub> evolution, Korean pine forest, Regression equation

## Introduction

CO<sub>2</sub> evolution from soil is one of the major processes in global carbon cycling. Increase amount of CO<sub>2</sub> in atmosphere is partly resulted from the disturbance of stabilized vegetation and land used. Some general information for forest ecosystem is available on soil respiration (Schlenter and Cleve 1985). Management practice in forestry clearly has significant effect on the magnitude of soil CO<sub>2</sub> evolution (Ewel *et al.* 1987). Comparison results of soil CO<sub>2</sub> evolution rates at sites in different conditions have been well gotten. But there are few papers about the relative magnitude of biomass of roots, litter and soil organic matter (Hendrickson *et al.* 1984). Site conditions should be selected specifically. The situations on soil CO<sub>2</sub> evolution in ecosystem of Korean pine natural forest is investigated in this study at Changbai Mountain of Northeast China.

## Methods

### Study site

The study site was selected in forest area with Korean pine (*Pinus koraiensis*) as dominant species. The site is located in the experimental plots adjacent to Forest Ecosystem Research Station in Changbai Mountain (42°24' N, 128°6' E) at the elevation of 760m. The stand is a mature Korean pine and broad-leaved tree mixed forest. Major species of broad-leaved trees are *Tilia amurensis*, *Fraxinus mandshurica*, *Quercus mongolica*, *Acer mono* and *Betula platyphylla*. The shrub species mainly consisted of *Acer tegmentosum*, *Ribes mandshurica*, *Deutzia amurensis*, *Corylus mandshurica* and *Lonicera chrysentha*. The herbaceous coverage was low with few species such as *Oxalis acetosella*, *Lilium dahuricum* and *Urtica angustifolia*. The dark brown forest soil under dense canopy was well for undisturbed thick litter and humus layer.

### Measurement of soil CO<sub>2</sub> evolution

Rate of soil CO<sub>2</sub> evolution was measured using an open airflow system connected with a bottomless glass chamber. The chamber was inserted into soil 10 cm deep. The remaining part over soil surface has 1.5 L in volume. The covered surface area was 154m<sup>2</sup> for each of the two chambers. The airflow through chamber by two pumps connected to inlet and outlet started immediately when soil surface was covered. The flow rate at inlet and outlet was the same as 1.2 L per minute, so that the positive or negative air pressure within chamber was avoided. CO<sub>2</sub> from soil respiration was well mixed with the flowing air by a mini-fan set in chamber. Sensors for measuring soil temperature were inserted into the soil of 5 cm deep inside and outside the covering areas. There was no significant difference in soil temperature inside and outside the chamber. When airflow started through chamber after 24h, gas samples were collected from a tube at the outlet every 2h using special bags made of polyene-aluminium film. Each of 3 parallel bags contained 1 L gas. At the same time, the contrast air samples were taken from atmosphere near the inlet tube. CO<sub>2</sub> concentrations of air samples were detected using an infrared gas analyzer soon, after the samples were taken and sent to the laboratory. Soil CO<sub>2</sub> evolution rate was calculated using the equation on CO<sub>2</sub> gas exchange (Sestak *et al.* 1971).

$$C_R = \frac{P_c \cdot F \cdot 10^{-6}}{A} \times \frac{44000}{22.4} \times \frac{273}{273+T} \times \frac{P_1}{P_0}$$

Where:

$C_R$  is rate of CO<sub>2</sub> evolution, mg·m<sup>-2</sup>·h<sup>-1</sup>;

$P_c$  is difference in CO<sub>2</sub> (ppm) between samples from inlet and outlet;

$F$  is air flow rate through chamber, L·h<sup>-1</sup>;

<sup>1</sup> This project was funded by the Opened Research Station of Changbai Mountain Forest Ecosystem, Chinese Academy of Sciences.

$A$  is soil surface area covered, m<sup>2</sup>;  
 $T$  is air temperature during measurement, °C;  
 $P_1$  is atmospheric pressure at measuring site, mb;  
 $P_0$  is atmospheric pressure at standard condition.

## Results and discussion

### Diurnal fluctuation of CO<sub>2</sub> evolution rate

Air samples out of the chambers were collected at 2 h interval from 6:00 to midnight in succession in June and August respectively. At each collection, 6 samples from chamber outlet and 3 samples from ambient atmosphere were taken for detecting CO<sub>2</sub> concentration. The values in Fig. 1 were average values from duplicate samples of two chambers.

At 6:00 in the morning, values both in June and August were comparatively low and were 220 and 280 mg/(m<sup>2</sup>•h)

respectively. The evolution rate gradually increased during a day. The higher values appeared at 18:00 and reached to 460 or 490 mg/(m<sup>2</sup>•h) respectively.

Growing season in this region was known as from April to October and summer time was from June to August. Compared to the value (0.7 g/(m<sup>2</sup>•h) by Gordon *et al* (1987) in spruce forest in mid-summer, our results were apparently lower. The highest value in Changbai Mountain area would be expected to appear in July in the region.

### Relationship between CO<sub>2</sub> evolution rate and soil temperature

The lowest value of soil temperature was 13 °C and appeared at 6:00 and the highest was 16.2 °C at 18:00. Diurnal change in CO<sub>2</sub> evolution rate was coincidentally with the change in temperature (Table 1).

Table 1. Values of CO<sub>2</sub> evolution and soil temperature in June

	Time									
	6:00	8:00	10:00	12:00	14:00	16:00	18:00	20:00	22:00	
Soil temperature /°C	13.0	13.8	14.5	15.0	15.7	16.0	16.1	15.2	14.8	
CO <sub>2</sub> evolution /mg•m <sup>-2</sup> •h <sup>-1</sup>	220	240	290	320	390	440	460	360	300	

According to the figures in Table 1, a regression equation was constructed as

$$Y=0.0812e^{0.2476x}$$

Where:  $Y$  is CO<sub>2</sub> evolution rate,  $X$  is temperature.




Fig.1. Diurnal fluctuations of CO<sub>2</sub> evolution rate in June

The relationship between evolution rate and soil temperature was found as an exponential form (Fig.2).

As the major factors affecting soil CO<sub>2</sub> evolution, the

magnitude of root biomass, organic matter, litter and soil animals, are also important. These factors could affect the of CO<sub>2</sub> evolution in different soil conditions at different sites. Our measurements indicated that the dynamic course of CO<sub>2</sub> evolution for a certain site apparently followed the change of soil temperature with the time course. For estimating the magnitude of soil CO<sub>2</sub> evolution with change of time, it is necessary to understand such dynamic change of different soil CO<sub>2</sub> evolution and to get soil temperature record of the sites.

### Estimation of CO<sub>2</sub> evolution from Korean pine forest soil

According to the regression equation on the relationship between CO<sub>2</sub> evolution rate, and the measured monthly average value of soil temperature, we estimated the monthly magnitude of CO<sub>2</sub> evolution in the growing season. The growing season in this region has about 183 d from mid-April to mid-October having detectable soil respiration. From Table 2, a total weight of CO<sub>2</sub> evolution was calculated for 10.4 ton per hectare during the growing season of the year.

Table 2. A calculation of soil CO<sub>2</sub> evolution in growing season

Month	April	May	June	July	August	September	October	Total
Average temperature /°C	4.50	7.80	14.60	16.90	15.70	11.20	5.10	
Evolution rate /mg•m <sup>-2</sup> •h <sup>-1</sup>	7	30	290	540	420	120	13	
CO <sub>2</sub> evolution daily /g•m <sup>-2</sup> •d <sup>-1</sup>	0.17	0.72	6.96	12.96	10.08	2.88	0.31	
CO <sub>2</sub> evolution monthly /g•m <sup>-2</sup> •month <sup>-1</sup>	2.55	22.32	208.80	401.76	312.48	86.40	9.61	1043.9
Total amount /kg •hm <sup>-2</sup> •a <sup>-1</sup>								10439

In a report by Ewel *et al.* (1987), an average value of carbon release rate from slash pine (*Pinus elliotii*) plantations in Florida was given as  $10.6 \text{ t}/(\text{hm}^2\cdot\text{a})$ . This figure was much higher than our value on the basis of carbon weight. A value of  $10.7 \text{ t}/(\text{hm}^2\cdot\text{a})$  was reported by Edwards and Harris (1977) for tennessee *Liriodendron tulipifera* forest. There was the greater difference for our measurement value on the basis of carbon ( $2.81 \text{ t}/(\text{hm}^2\cdot\text{a})$ ). And those values might be attributed to the difference of the latitudes and site conditions. The dense canopy and lower soil temperature were found to be the major reasons on our lower value in carbon release in Korean pine forest.




Fig. 2. Relationship between  $\text{CO}_2$  evolution rate and soil temperature measured in June under the dense canopy of Korean pine natural forest

### Acknowledgments

The authors would like to express their sincere thanks to Mr. Lin Jihui for his assistance in field measurements and

to Ms. Zhao Jiewei for manuscript preparation.

### References

- Edwards, N. T. and Harris, W. F. 1997. Carbon cycling in a mixed deciduous forest floor. *Ecology*. 58: 431-437.
- Ewel, K. C., Cropper, W. P. J. and Gholz, H. L. 1987. Soil  $\text{CO}_2$  evolution in Florida slash pine plantations. I. Changes through time. *Can. J. For Res.* 17: 325-329.
- Gordon, A. M., Schlentner, R. E. and Van Cleve, K. 1987. Seasonal patterns of soil respiration and  $\text{CO}_2$  evolution following harvesting in the white spruce forests of interior Alaska. *Can. J. For. Res.* 17: 304-310.
- Hendrickson, O. Q. and Robinson, J. B. 1984. Effects of roots and litter on mineralization processes in forest soil. *Plant Soil*. 80: 391-405.
- Schlentner, R. E. and Van Cleve, K. 1985. Relationships between  $\text{CO}_2$  evolution from soil, substrate temperature, and substrate moisture in four mature forest types in interior Alaska. *Can. J. For. Res.* 15: 97-107.
- Sestak, Z., Catsky, J. and Jarvis, P. G. 1971. Plant photosynthetic production-Manual of Methods. Dr. W. Junk, N. V. Publishers, The Hague. pp. 53-163.
- Van Cleve, K. and Moore, T. A. 1978. Cumulative effects of nitrogen, phosphorus, and potassium fertilizer additions on soil respiration, pH, and organic matter content. *Soil Sci. Soc. Am. J.* 42: 121-124.

(Responsible editor: Zhu Hong)